

# HAAD-RN<sup>Q&As</sup>

HAAD Licensure Examination for Registered Nurses

**Pass HAAD HAAD-RN Exam with 100% Guarantee**

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.pass2lead.com/haad-rn.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by HAAD  
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



**QUESTION 1**

Salem has Alzheimer's disease. He is agitated and repeatedly asks to go home. The most appropriate nursing intervention for him is to:

- A. Isolate him in a single room
- B. Find activities to keep him occupied
- C. Ask the physician to discharge him
- D. Administer a minor tranquilizer

Correct Answer: B

---

**QUESTION 2**

To maintain the airway and promote respiratory function, the best position for the nurse to place the unconscious patient in the Post Anesthesia Care Unit (PACU) is:

- A. Supine
- B. Lateral
- C. Trendelenberg
- D. Fowler's

Correct Answer: B

---

**QUESTION 3**

The patient has a nursing diagnosis of altered cerebral tissue perfusion related to cerebral edema. An appropriate nursing intervention for this problem is to:

- A. Elevate the head of the bed 30 degrees
- B. Provide a position of comfort with knee flexion
- C. Provide uninterrupted periods of rest
- D. Ensure adequate hydration with mannitol

Correct Answer: A

---

**QUESTION 4**

A patient complains of pain in his right arm. The physician orders codeine 45 mg and aspirin 650 mgs every 4 hours PRN. Each codeine tablet contains 15mg of codeine and each aspirin tablet contains 325mg of aspirin. What should the

nurse administer?

- A. 2 codeine tablets and 4 aspirin tablets
- B. 4 codeine tablets and 3 aspirin tablets
- C. 3 codeine tablets and 2 aspirin tablets
- D. 3 codeine tablets and 3 aspirin tablets

Correct Answer: C

---

**QUESTION 5**

Which of the following tasks requires specialized education and should be performed by the nurse only after the training has been completed?

- A. Administering a dose of promethazine (Phenergan) via intravenous push (IVP)
- B. Applying a transdermal fentanyl (Duragesic)
- C. Instilling tobramycin (Tobrex) ophthalmic solution
- D. Beginning an intravenous infusion of cyclophosphamide (Cytosan)

Correct Answer: D

---

**QUESTION 6**

Symptoms of alcohol withdrawal include:

- A. Euphoria, hyperactivity and insomnia
- B. Depression, suicidal ideation and hypersomnia
- C. Diaphoresis, nausea and vomiting and tremors
- D. Unsteady gait, nystagmus and profound disorientation

Correct Answer: C

---

**QUESTION 7**

The minimal amount of urine that a post-operative patient weighing 60 kgs should pass is?

- A. 120 ml/hr
- B. 90 ml/hr
- C. 60 ml/hr

D. 30 ml/hr

Correct Answer: D

---

**QUESTION 8**

To prevent post-operative thrombophlebitis, which one of the following measures is effective?

- A. Elevation of the leg on two pillows
- B. Using of compression stocking at night
- C. Massage the calf muscle frequently
- D. Performing leg exercises

Correct Answer: D

---

**QUESTION 9**

Which of the following techniques should the nurse implement to prevent the patient's mucous membranes from drying when the oxygen flow rate is higher than 4 liters per minute?

- A. Use a non rebreather mask
- B. Add humidity to the delivery system
- C. Use a high flow oxygen delivery system
- D. Ensure that the prongs are in the nares correctly

Correct Answer: B

---

**QUESTION 10**

The nurse knows that the greatest risk for a patient with a ruptured ectopic pregnancy is:

- A. Hemorrhage leading to hypovolemic shock
- B. Strictures and scarring of the fallopian tube
- C. Adhesions and scarring from blood in the abdomen
- D. Infertility resulting from treatment with a salpingectomy

Correct Answer: A

---

**QUESTION 11**

A patient has had a total hip joint replacement. Which of the following actions should the nurse consider for the patient's daily recommended exercise program?

- A. Administering an analgesic before exercising
- B. Discontinuing the program if the patient dislikes it
- C. Continuing exercises in spite of severe pain
- D. Evaluating effectiveness of exercise based on pain scale

Correct Answer: A

---

#### QUESTION 12

The purpose of a cardiac pacemaker is to:

- A. Initiate and maintain the heart rate when SA node is unable to do so
- B. Stabilize the heart rate when it is above 100 beats per minute
- C. Stabilize the heart when the patient has had a heart attack
- D. Regulate the heart when the patient is going for open heart surgery

Correct Answer: A

---

#### QUESTION 13

When caring for a patient with hepatic encephalopathy the nurse may carry out the following orders: give enemas, provide a low protein diet, and limit physical activities.

These measures are performed to:

- A. Minimize edema
- B. Decrease portal pressure
- C. Reduce hyperkalemia
- D. Decrease serum ammonia

Correct Answer: D

---

#### QUESTION 14

Order: 1000 ml of D5W to run for 12 hours. Using a micro drip set calculate the drops per minute (gtts/ min):

- A. 20 gtts/min

B. 45 gtts/min

C. 60 gtts/min

D. 83 gtts/min

Correct Answer: D

---

**QUESTION 15**

To promote accuracy of self-monitoring blood glucose by patients the nurse should:

A. Retrain patients periodically

B. Direct patients to rotate testing sites

C. Advise patients to buy new strips routinely

D. Compare results from patient's meter against lab results

Correct Answer: A

[Latest HAAD-RN Dumps](#)

[HAAD-RN Study Guide](#)

[HAAD-RN Exam Questions](#)